



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

DIOPHANTINE ANALYSIS.

127. Proposed by F. P. MATZ, Sc. D., Ph. D., Reading, Pa.

Can there be determined three cube numbers whose sum is the product of two squares?

AVERAGE AND PROBABILITY.

165. Proposed by HENRY HEATON, Atlantic, Iowa.

What is the average length of all straight lines that can be drawn within a given square parallel to one of the diagonals?

166. Proposed by F. P. MATZ, Sc. D., Ph. D., Reading, Pa.

Find the average area intercepted by two non-intersecting chords drawn at random in a given circle.

167. Proposed by R. D. CARMICHAEL, Hartselle, Ala.

A line l is divided into n segments by $n-1$ points taken at random on it; find the mean value of the product of p of the segments, the p segments being taken at random and p being less than n .

GROUP THEORY.

8. Proposed by L. E. DICKSON, Ph. D., The University of Chicago.

In a chess tournament between eight players, there are seven rounds, the eight players being paired in each round, every pair to be matched once and but once in the tournament. List the possible programs different except as to notation, *i. e.*, not transformable into each other by a substitution on eight letters. Give the number of conjugate programs of each representative retained.

MISCELLANEOUS.

149. Proposed by F. P. MATZ, Ph. D., Sc. D., Reading, Pa.

Given $\sin^{-1}u + \sin^{-1}\frac{1}{2}u = \frac{1}{2}\pi$, to find u .